

Name _____

Practice Calculations Test Problems

Do-overs Allowed - only 3 significant figures in answers needed.

In the following formulas, use $G=6.67\times 10^{-11}\frac{\text{N}\cdot\text{m}^2}{\text{kg}^2}$ and $c=3.00\times 10^8\text{ m/s}$.

1. Calculate the gravitational force in newtons using the formula $F=G\frac{m_1m_2}{d^2}$ where,
 $m_1=5.3\times 10^{23}\text{kg}$, $m_2=64.9\text{ kg}$ and $d=8.3\times 10^8\text{ m}$.

2. Calculate the dimensionless relativistic factor $\gamma=\frac{1}{\sqrt{1-\frac{v^2}{c^2}}}$ where $v=2.8\times 10^8\text{ m/s}$.

3. Using $E=mc^2$, calculate the energy in joules released when $3.7\times 10^{-3}\text{ kg}$ is converted to energy.

4. Using $\lambda=\frac{c}{f}$, calculate the wavelength in meters of a photon that has a frequency of $5\times 10^{15}\text{ Hz}$.

5. Acetic acid, the active ingredient in vinegar, has a pH of 4.756. Calculate the concentration in mol/L of hydronium ions in vinegar using $[\text{H}_3\text{O}^+]=10^{-\text{pH}}$.

6. The concentration of hydronium ions in household ammonia is $2.34\times 10^{-12}\text{ mol/L}$. Calculate the pH of household ammonia using $\text{pH}=-\log_{10}[\text{H}_3\text{O}^+]$

Practice Calculations Test Problems - Solutions
Do-overs Allowed - only 3 significant figures in answers needed.

In the following formulas, use $G=6.67 \times 10^{-11} \frac{\text{N} \cdot \text{m}^2}{\text{kg}^2}$ and $c=3.00 \times 10^8 \text{ m/s}$.

1. Calculate the gravitational force in newtons using the formula $F=G \frac{m_1 m_2}{d^2}$ where,
 $m_1=5.3 \times 10^{23} \text{ kg}$, $m_2=64.9 \text{ kg}$ and $d=8.3 \times 10^8 \text{ m}$.

$$F=(6.67 \times 10^{-11} \frac{\text{N} \cdot \text{m}^2}{\text{kg}^2}) \frac{(5.3 \times 10^{23} \text{ kg}) \cdot (64.9 \text{ kg})}{(8.3 \times 10^8 \text{ m})^2} = 3.330 \times 10^{-3} \text{ N} = 3.330 \text{ mN}$$

2. Calculate the dimensionless relativistic factor $\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$ where $v=2.8 \times 10^8 \text{ m/s}$.

$$\gamma = \frac{1}{\sqrt{1 - \frac{(2.8 \times 10^8 \text{ m/s})^2}{(3.00 \times 10^8 \text{ m/s})^2}}} = \frac{1}{\sqrt{1 - \left(\frac{2.8 \times 10^8 \text{ m/s}}{3.00 \times 10^8 \text{ m/s}}\right)^2}} = 2.785$$

3. Using $E=mc^2$, calculate the energy in joules released when $3.7 \times 10^{-3} \text{ kg}$ is converted to energy.

$$E=(3.7 \times 10^{-3} \text{ kg}) \cdot (3.00 \times 10^8)^2 = 3.33 \times 10^{14} \frac{\text{kg} \cdot \text{m}^2}{\text{s}^2} = 3.330 \times 10^{14} \text{ J} = 3.33 \times 10^{14} \text{ N} \cdot \text{m} = 3.33 \times 10^{14} \text{ J} = 333 \text{ TJ}$$

4. Using $\lambda = \frac{c}{f}$, calculate the wavelength in meters of a photon that has a frequency of $5 \times 10^{15} \text{ Hz}$.

$$\lambda = \frac{3.00 \times 10^8 \text{ m/s}}{5 \times 10^{15} \text{ Hz}} = 6.00 \times 10^{-8} \text{ m} = 0.0600 \mu\text{m} = 60.0 \text{ nm}$$

5. Acetic acid, the active ingredient in vinegar, has a pH of 4.756. Calculate the concentration in mol/L of hydronium ions in vinegar using $[\text{H}_3\text{O}^+] = 10^{-\text{pH}}$.

$$[\text{H}_3\text{O}^+] = 10^{-\text{pH}} = 10^{-4.756} = 1.754 \times 10^{-5} \text{ mol/L}$$

6. The concentration of hydronium ions in household ammonia is $[\text{H}_3\text{O}^+] = 2.34 \times 10^{-12} \text{ mol/L}$. Calculate the pH of household ammonia using $\text{pH} = -\log_{10}[\text{H}_3\text{O}^+]$

$$\text{pH} = -\log_{10}[\text{H}_3\text{O}^+] = -\log_{10}(2.34 \times 10^{-12} \text{ mol/L}) = 11.63$$

Note: pH has no units, but must be calculated from $[\text{H}_3\text{O}^+]$ with units of mol/L .